Location of Clinic	In Charge of	Clinic			
		Opened	Closed	Inoculated	Totals
City of Fresno	City H. O.	September 29, 1924	November 28, 1924	20,812	
Private offices	Dr. C. Mathewson Physicians of	September-October	November, 1924	19,849	
Fresno County	Fresno County H. O. Dr. G. C. Long	October 20	December 31	22,230	
Private offices	Physicians of County	October-November	December	4,671	
	Total inoculated wit	hin Fresno County			67,562
Tulare County Kings County Madera County Merced County *Sacramento County	Dr. G. J. Telfer Dr. G. J. Telfer Dr. G. J. Telfer Dr. G. J. Telfer Dr. W. W. Cress	October and November October and November October and November October and November November Total in other counties		14,034 8,208 5,088 5,000 2,730	
				I	35,060
		Total inoculations		102.622	

of an inch in diameter were denuded, extending down into the Malpigian layer of the derma, and very seldom resulted in any bleeding. Fresh vaccine was applied to these denuded areas and the wounds immediately covered with two or three layers of aseptic gauze, held in place by adhesive plaster. The reaction was observed on the eighth day and fresh gauze applied. No serious results were observed when this method was followed. Many persons who were successfully inoculated by this method were hard to convince that they had been successfully immunized. "But my arm never hurt me" was heard innumerable times.

The relation of immunization to protection against smallpox bore out the evidence that history has repeatedly demonstrated since the days of Jenner.

	Recoveries	Deaths
History unknown	1	5
Never successfully immunized		16
Successfully immunized under five		••••
Successfully immunized five to ten		••••
Successfully immunized over ten ye	ars 9	4
Totala	1/5	95

Many cases were seen where a typical vaccinia developed at the same time that an unmodified smallpox rash grew, indicating that the unfortunate victim had been inoculated just two or three days too late after having been infected with smallpox. So far as observed, these late inoculations did not seem to have any influence upon the severity of the attack. All who had passed through the maximum stage of their vaccine inoculation before being exposed, or before the incubation period of smallpox had passed after having been exposed, were protected against the disease and did not get it. This amply demonstrated the statement that "an immune person cannot develop smallpox." Immunity was occasionally found to have been shortened in certain individuals to a few years, probably dependent on some peculiar systemic conditions possessed by these particular persons; however, I never found immunity to have been shortened to less than three years.

Yours very truly,

Allen F. Gillihan, M. D., County Health Officer.

Subject of Following Letter: Cinch Shortening Operation for Strabismus

To the Editor:—It has recently come to my attention that a story is being circulated to the effect that I have been guilty of a violation of medical ethics in trying to keep secret my cinch shortening operation

for the correction of strabismus and other troubles of the ocular muscles.

I hope you will publish this statement, as I know of no other way to get the facts to all who may have heard the story.

The operation was devised by me in 1911 and reported, after my first operation, in the Journal of the American Medical Association for March 2, 1912. I was then in the Army Medical Corps, which afforded but little opportunity to see much muscle work. As stated in the paper, the report was made at once in hopes of interesting some who saw a great deal of that kind of work.

My second paper was in the Archives of Ophthalmology for 1914, and later in the same year, at the request of Dr. Casey Wood, I took part in a symposium on the ocular muscles published in the Ophthalmic Record. Also, at his request, I furnished a condensed description for the Encyclopedia of Ophthalmology. In 1915 a short paper, with demonstration, was given at the meeting of the Pacific Coast Oto-Ophthalmological Society. The next paper was at the 1916 state medical meeting at Fresno.

The operation was brought up to date by reporting forty-two operations at the 1916 meeting of the American Medical Association at Detroit. While on that trip I operated in Chicago for Dr. Casey Wood, and at the Illinois Eye and Ear for Doctor Orcutt.

Since that year I have read many papers, and at all meetings have been busy demonstrating the principle of the method to anyone interested.

I resigned from the Army in 1914 and started practice in Oakland in 1915. During that year several ophthalmologists brought patients to me and assisted at the operations.

Operations were done at the Knapp Eye Hospital and the Wills Eye Hospital in 1914. In 1930, operations were done for Doctor Orcutt in Chicago during the Academy meeting. In 1931, eight squints were operated at Salt Lake City for the Utah State Medical Society. In November, 1931, a number of operations were done for the Southern Medical Association meeting in New Orleans.

In 1924 Dr. Joseph L. McCool, then of Portland, made a week's visit and assisted at eight or ten operations I had saved up for that purpose, and last December Dr. Donald O'Rourke of Denver made a similar visit for the same purpose.

In view of all these provable facts, I think it is quite apparent that the story of my withholding information concerning the technique of the operation is without foundation in fact.

RODERIC O'CONNOR.

450 Sutter Street, San Francisco.